

Technical Data

Acriflavin-Cefsulodin-Vancomycin Supplement (ACV Supplement)

FD284

Recommended for the isolation of Enterohemorrhagic E. coli (EHEC).

Composition

Per vial sufficient for 500 ml medium

*Ingredients Concentration
Acriflavin 50mg
Cefsulodin 40mg
Vancomycin 50mg

Directions:

Rehydrate the contents of 1 vial aseptically with 5 ml of sterile distilled water. Mix well and aseptically add to 500 ml of sterile, cooled (45-50 $^{\circ}$ C) Buffered Peptone Water w/Pyruvate M1851 . Mix well and dispense as desired.

Storage and Shelf Life

Store at 2 - 8°C. Use before expiry date on the label.

Revision: 1 / 2012

* Not For Medicinal Use

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.