

S. aureus recombinant LukD

Catalog #: 0530-003

Lot #: 1508003

Description: Purified, *Staphylococcus aureus* recombinant leukocidin-D (rLukD). The rLukD (tag free) is expressed in *E. coli* and purified by FPLC. The theoretical molecular weight of the protein is 36,889 Daltons.

Storage: 2-3 weeks at -20°C, -80°C long term

Size: 100 µg of protein is supplied in PBS at a concentration of **1.085 mg/mL**. Protein demonstrates a molecular weight of approximately 37 kDa.

Endotoxin: 28.713 Endotoxin units/mg

Relevance: This protein may be used in functional toxicity assays in combination with rLukE, or as a control protein in ELISA assays or Western blotting when detecting toxins produced by different strains of *S. aureus*.

Recommended Dilutions:


ELISA: Assay-dependent dilution.

WB: Assay-dependent dilution; internal QC demonstrates detection of 100 ng of rLukD protein using anti-PVL LukF polyclonal antibody (cat# 0312-002) in Western blotting.


Cytotoxicity assay: Cytotoxicity can be detected in human neutrophils when used in combination with LukE.

For additional *S. aureus* products, please visit:
<http://ibtbioservices.com/index.php/product-and-reagents/staphylococcal-products>

SDS-PAGE and Western Blot Detection



(A) SDS-PAGE of rLukD: 1 µg (lane 1) and 5 µg (lane 2). (B) Western blot detection of rLukD at 100 ng (lane 3), using IBT's anti-LukF-PV polyclonal antibody (cat# 0312-002) at 0.5 µg/mL and an anti-rabbit IgG-HRP conjugate followed by substrate.



Toxin Functionality: Human promyelocytic leukemia cell line HL60 was differentiated into neutrophils by treatment with DMSO. Neutrophils were incubated with serial dilutions of rLukE and rLukD at equimolar concentration for 3 hours at 37°C with 5% CO₂ and 95% humidity. Cellular viability was determined by adding XTT and incubation for additional 16 hours. Cells were centrifuged and the OD determined in the supernatants at 470/690 nm. Red circles represents the current lot 1508003 and blue circles represents the previous lot 1411005. EC₅₀ were found to be 19.4 nM for the current lot and 24.1 nM for the previous lot.

Intended for research use only. Not for human, therapeutic, or diagnostic applications.

The buyer cannot sell or otherwise transfer this product for Commercial Purposes without written approval of Integrated BioTherapeutics, Inc.
Copyright 2015. Integrated BioTherapeutics, Inc. All rights reserved.