

Technical Data

HiEncapTM Buffered Peptone Water

EC1494ID

HiEncapTM Buffered Peptone Water is used as a pre-enrichment medium for increasing the recovery of injured Salmonella species from foods prior to selective enrichment and isolation. The composition and performance criteria of this medium are as per the applications laid down in ISO 6579-2002.

Composition**

Ingredients	Gms / Litre
Enzymatic Digest of Casein	10.000
Sodium chloride	5.000
Disodium hydrogen phosphate,12H ₂ O	9.000
Potassium dihydrogen phosphate	1.500
Final pH (at 25°C)	7.0 ± 0.2

^{**}Formula adjusted, standardized to suit performance parameters

Directions

Each capsule contains 10.04 gms of dehydrated medium. Suspend 1 capsule in 500ml (2 capsules in 1000 ml) distilled or purified water. Heat to boiling to dissolve the medium completely. Dispense as desired and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Principle And Interpretation

Microorganisms that are subjected to environmental stresses may become structurally or metabolically damaged or injured. These microorganisms are unable to replicate in selective environments. Therefore these injured organisms must be resuscitated or permitted to repair the damage by incubation in an appropriate, non-selective environment (1). Edel and Kampelmacher (2) noted that sublethal injury to Salmonellae may occur in many food preservation processes. Enriching injured cells in Lactose Broth (pH 6.9) may be further detrimental to their recovery (3). Pre-enrichment in Buffered Peptone Water (M1494I) at 35°C for 18-24 hours results in repair of injured cells (4). The buffering system prevents bacterial damage due to change in the pH of the medium. Recently ISO committee has also recommended this pre-enrichment medium for the detection of *Enterobacteriaceae* from food stuffs and other materials (5).

Inoculate 10 grams specimen in 50 ml of Buffered Peptone Water (EC1494ID) and incubate at 35°C for 18 hours. Transfer 10 ml from this medium to 100 ml of Mueller Kauffman Tetrathionate Novobiocin Broth Base (M1496I) and Rappaport Vassiliadis Soya Broth (RVS Broth) (M1491) and incubate at 43°C for 24-48 hours and then subculture on selective media like XLD Agar, Modified (M031I). Examine the plates for colonies of *Salmonella* species.

Quality Control

Appearance

Gelatin capsule containing, cream to yellow granular media

Colour and Clarity of prepared medium

Light yellow coloured clear solution without any precipitate

Reaction

Reaction of 2.0% w/v aqueous solution at 25°C. pH : 7.0 ± 0.2

pН

6.80-7.20

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.(Recovery is observed on XLD Agar, M031I)

Cultural Response

Organism Inoculum Growth Recovery (CFU)

HiMedia Laboratories Technical Data

Cultural Response			
Salmonella Enteritidis ATC	C 50-100	luxuriant	>=50%
13076			
Salmonella Typhi ATCC	50-100	luxuriant	>=50%
6539			
Salmonella Typhimurium	50-100	luxuriant	>=50%
ATCC 14028			
Escherichia coli ATCC	50-100	luxuriant	>=50%
25922			
Pseudomonas aeruginosa	50-100	luxuriant	>=50%
ATCC 27853			

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

Disclaimer:

- 1. Vanderzant C. and Splittstoesser D. F., (Eds.), 1992, Compendium of Methods for the Microbiological Examination of Foods, 3rd Ed., APHA, Washington, D.C.
- 2. Edel W. and Kampelmacher E. H., 1973, Bull. Wld. Hlth. Org., 48: 167.
- 3. Angelotti R., 1963, "Microbiological Quality of Foods", Academic Press, New York.
- 4. Sadovski A. Y., 1977, J. Food Technol., 12.85.
- 5. International Organization for Standardization (ISO), 2002, Draft ISO/DIS, 6579.

Revision: 00 / 2014

(€

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.