



Bile Esculin Azide Agar

M493

Bile Esculin Azide Agar is a selective medium used for isolation and presumptive identification of faecal Streptococci.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	17.000
Beef extract	5.000
Proteose peptone	3.000
Oxgall	10.000
Esculin	1.000
Ferric ammonium citrate	0.500
Sodium chloride	5.000
Sodium azide	0.150
Agar	15.000
Final pH (at 25°C)	7.1±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 56.65 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Caution: Sodium azide has a tendency to form explosive metal azides with plumbing materials. It is advisable to use enough water to flush off the disposables.

Principle And Interpretation

Group D Streptococci possess the group D lipoteichoic acid antigen in their cell walls. Former Group D species, which are predominant normal inhabitants of the human gastrointestinal tract, are termed as faecal Streptococci or Enterococci (1). The unique ability of Enterococci to split esculin was reported by Meyer and Schonfeld (2). Enterococci and Group D Streptococci hydrolyse esculin to esculetin and dextrose, which reacts with ferric citrate producing brownish black precipitate (3). The use of esculin hydrolysis in identification of Enterococci was first cited by Rochaix (4). Bile Esculin Agar was originally formulated by Swan (6) for the isolation and identification of Group D Streptococci from food. Facklam and Moody (7, 8) further reported that using Bile Esculin Agar, Group D Streptococci could be differentiated from non Group D Streptococci. Bile Esculin Agar was also shown to aid differentiation of *Enterobacteriaceae*, *Klebsiella*, *Enterobacter*, *Serratia* from other Enterobacteriaceae genera (9) on the basis of esculin hydrolysis. However, other tests such as salt tolerance should be performed for identifying Enterococci (5).

Bile Esculin Azide Agar is a modification of Bile Esculin Agar (6, 8) as per Isenberg (10). In this medium the bile concentration is reduced and additional sodium azide is incorporated.

Casein enzymic hydrolysate, proteose peptone and beef extract serves as sources of carbon, nitrogen, amino acids, vitamins and essential growth nutrients. Oxgall and sodium azide inhibits most of the other accompanying bacteria. Esculin in the medium is hydrolyzed to esculetin and dextrose. Esculetin reacts with ferric citrate to form a dark brown or black complex, visualized as a zone of black precipitate around the colonies. If the media is dispensed in tubes in the form of slants, a positive reaction is indicated by blackening of more than half of the slant within 24-48 hours. If blackening is totally absent or if less than half of the slant is blackened within 24-48 hours, the test is negative. Viridans Streptococci sometimes exhibit a weak positive reaction. Also, *Leuconostoc*, *Pediococcus*, *Lactococcus* species causing human infections give a positive bile esculin test (11). To enhance the growth of Enterococci, Bile Esculin Agar can be supplemented with 50ml/l horse serum (3).

Suspected water samples are filtered using membrane filters. These membrane filters are aseptically placed on Slanetz and Bartely Medium (M612I). Red or maroon coloured colonies observed after incubation are further confirmed by aseptically

**Disclaimer :**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.