IMMOLASE™ DNA Polymerase

Shipping: Dry/Blue Ice Catalog numbers

BIO-21046: 250 Units

Batch No.: See vial BIO-21047: 500 Units

Concentration: 5 u/uL BIO-21048: 5000 Units



Store at -20°C

The IMMOLASE is shipped on dry/blue ice. On arrival store at -20 °C for optimum stability. Repeated freeze/thaw cycles should be avoided.

Expiry:

When stored under the recommended conditions and handled correctly, full activity of the kit is retained until the expiry date on the outer box label.

Safety precautions:

Storage and stability:

Please refer to the material safety data sheet for further information.

Unit definition:

One unit is defined as the amount of enzyme that incorporates 10nmoles of dNTPs into acid-insoluble form in 30 minutes at 72 $^{\circ}\text{C}.$

Notes:

This reagent has been manufactured under 13485 Quality Management System and is suitable for further manufacturing use as an IVD component.

Features

- Outstanding and robust performance
- Excellent specificity
- Convenient set up at room temperature
- Available in ready-to-go versions ImmoMix™ and ImmoMix™ Red

Applications

- Hot-start PCR assays
- TA cloning
- Ultra-high specificity for multiplex reactions
- Low-copy number templates

Description

IMMOLASE™ DNA Polymerase is a heat-activated thermostable DNA polymerase isolated from a novel organism. IMMOLASE provides improved specificity as compared to standard polymerases and can eliminate the presence of non-specifics, such as primer-dimers and mis-primed products. IMMOLASE is inactive at room temperature and therefore prior to PCR cycling, requires activation by heat treatment for 10 minutes. Subsequently, the reaction can be handled according to the user's existing protocols for thermostable DNA Polymerases.

Components

Product	250 Units	500 Units	5000 Units
IMMOLASE DNA Polymerase	1 x 50 μL	1 x 100 μL	10 x 100 μL
10x ImmoBuffer	1.2 mL	2 x 1.2 mL	20 x 1.2 mL
50 mM MgCl ₂ Solution	1.2 mL	1 x 1.2 mL	10 x 1.2 mL

Associated Products

Product	Pack size	Cat. No.
dNTP Set	4 x 25 µmol	BIO-39025
dNTP Mix	500 μL	BIO-39028
ImmoMix™	500 Reactions	BIO-25020
ImmoMix [™] Red	100 Reactions	BIO-25021
Agarose	100 g	BIO-41026

Product Citations:

- 1. Payne, B.A. Nature Gene. 43, 806-810 (2011)
- 2. Massire, C., et al. J. Clin. Microbiol. 49, 908 917 (2011)
- 3. Ashton, E.J. Meth. Mol. Biol. 688, 1-6 (2011)
- 4. Kaczmarek, K., et al. Mol. Biol. Cell 22, 1766 1779 (2011)
- 5. Scoville, A.G. & Pfender, M.E. PNAS 107(9), 4260-4263 (2010)

PCR Reaction conditions (for a 50 μL reaction)

 $10x \ ImmoBuffer \qquad \qquad 5 \ \mu L$ $100 \ mM \ dNTP \ Mix^* \qquad \qquad 0.5 \ \mu L$ $50 \ mM \ MgCl_2 \ Solution \qquad \qquad 3.0 \ \mu L$ $Template \ and \ primers \qquad \qquad As \ required$ $IMMOLASE \qquad \qquad 1 \ \mu L$

η μΕ

Water (ddH $_2$ 0) Up to 50 μ L

* Bioline 100 mM dNTP Mix is available as a separate product (BIO-39028)

Activation: pre-heating step at 95°C for 10 minutes

Denaturation: 94-96 °C

Annealing: depends on primer Tm

Extension: 72 °C (allowing 15-30 seconds/kb)

General Considerations:

Pre-incubate at 95 °C for 10 minutes. Subsequently, the reaction can be treated according to the user's existing protocols.

If extension time exceeds 2.5 minutes, a maximum of 30 cycles should be used. Increasing the number of cycles may lead to smearing when run on an agarose gel.

The ideal MgCl $_2$ concentration in the reaction is likely to be 1.5 - 2.5 mM (final concentration), but some optimization may be necessary to achieve the best possible results. For first tests, use no less than 1 unit of IMMOLASE in a 50 μ L reaction.

This data is intended for use as a guide only; conditions will vary from reaction to reaction and may need optimization.

Notes

IMMOLASE and ImmoMix are Trademarks of Bioline Reagents Ltd.

Bioline Reagents Ltd Bioline USA Inc. Bioline GmbH Bioline (Aust) Pty. Ltd Bioline France Meridian Bioscience Asia Pte Ltd UNITED KINGDOM **GERMANY AUSTRALIA** FRANCE SINGAPORE USA Tel: +44 (0)20 8830 5300 Tel: +1 508 880 8990 Tel: +49 (0)337 168 1229 Tel: +61 (0)2 9209 4180 Tel: +33 (0)1 42 56 04 40 Tel: +65 6774 7196 Fax: +44 (0)20 8452 2822 Fax: +1 508 880 8993 Fax: +61 (0)2 9209 4763 Fax: +49 (0)9 70 06 62 10 Fax: +65 6774 6441

Website: www.bioline.com/ email: info@bioline.com