

# **Technical Data**

# Eugonic LT 100 Broth Base w/o Tween 80

# M1517

Eugonic LT 100 Broth Base is recommended for the enrichment and detection of mesophilic aerobic bacteria present in cosmetic products. The composition and performance criteria of the medium are as per the specifications laid down in ISO 21149.

## Composition\*\*

Ingredients	Gms / Litre
Casein enzymic hydrolysate	15.000
Papaic digest of soyabean meal	5.000
Sodium chloride	4.000
L-Cystine	0.700
Sodium sulphite	0.200
Glucose	5.500
Egg lecithin	1.000
Tritox X-100	1.000
Final pH ( at 25°C)	7.0±0.2

\*\*Formula adjusted, standardized to suit performance parameters

# Directions

Suspend 32.4 grams in 1000 ml distilled water containing 5 grams of polysorbate 80(Tween 80). Heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

# **Principle And Interpretation**

Eugonic LT 100 Broth Base was developed by Pelczar and Vera (1) for cultivation of fastidious organisms like *Brucella* . Eugon media were developed to obtain eugonic (luxuriant) growth of fastidious microorganisms like *Brucella* which are otherwise difficult to cultivate (3). The unenriched medium supports rapid growth of lactobacilli associated with cured meat products, dairy products and other foods. Eugonic media is quite similar to Tryptone Soya Agar (M290) but more bacterial propagation is expected on Eugonic media. Organisms like *Bordetella* and *Neisseria* grow luxuriantly in Eugon Media because large amount of sulfur and carbon sources have been added in the formula. Eugonic LT 100 Broth Base can be used for growth of a variety of fastidious microorganisms like *Neisseria, Francisella* and *Brucella*. The composition of the medium is as per ISO (4)for the detection of mesophilic aerobic bacteria from cosmetic products.

Casein enzymic hydrolysate and papaic digest of soyabean meal provide the nitrogen, vitamins and amino acids, which supports the growth of fastidious microbial species. The high concentration of glucose is the energy source for rapid growth of bacteria. L-Cystine and sodium sulphite are added to stimulate growth. Sodium chloride maintains the osmotic balance of the media. The high carbohydrate content along with high sulfur (cystine) content improves growth with chromogenicity (2). Lecithin and polysorbate 80 in Eugonic LT 100 Medium w/o Tween 80 neutralize antimicrobial agents hence this medium can be used as a neutralizing diluent.

# **Quality Control**

#### Appearance

Cream to yellow homogeneous free flowing powder

## Colour and Clarity of prepared medium

Yellow coloured, clear to slightly opalescent solution.

## Reaction

Reaction of 3.24% w/v aqueous solution at 25°C. pH : 7.0±0.2

**pH** 6.80-7.20

**Cultural Response** 

Cultural characteristics observed after an incubation at 35-37°C for 24-48 hours (fungal cultures incubated at 25-30°C for 2-7 days).

<b>=</b> / <b>uuj</b> s).		
Cultural Response		
Organism	Inoculum (CFU)	Growth
Cultural Response		
Bacillus pumilus ATCC 14884	50-100	good
Candida albicans ATCC 26790	50-100	good
Lactobacillus fermentum ATCC 9338	50-100	good
Streptococcus pneumoniae ATCC 6303	50-100	good-luxuriant (under 3-5% CO2)
Streptococcus pyogenes ATCC 19615	50-100	good-luxuriant (under 3-5% CO2)
Staphylococcus aureus ATCC 25923	50-100	good-luxuriant
Staphylococcus aureus ATCC 6538	50-100	good
Bacillus subtilis ATCC 6633	50-100	good
Pseudomonas aeruginosa ATCC 9027	50-100	good
Escherichia coli ATCC 8739	50-100	good
Candida albicans ATCC 10231	50-100	good
Neisseria meningitidis ATCC 13090	250-100	good

#### **Storage and Shelf Life**

Store at below 30°C in tightly closed container and prepared media at 2-8°C. Use before expiry period on the label.

#### Reference

1.Pelczar and Vera J., 1949, Milk Plant Monthly 38:30

2.Frank H. A., 1955, J. Bacteriol., 70:269.

3.MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams & Wilkins, Baltimore, Md.

4.ISO 21149 (2006) Cosmetics-Microbiology- Enumeration and detection of aerobic mesophilic bacteria.

Revision : 1 / 2011

#### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia<sup>™</sup> publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia<sup>™</sup> Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

HiMedia Laboratories Pvt. Ltd. A-516, Swastik Disha Business Park, Via Vadhani Ind. Est., LBS Marg, Mumbai-400086, India. Customer care No.: 022-6147 1919 Email: techhelp@himedialabs.com Website: www.himedialabs.com

# CE