



Technical Data

HiEncap™ YT Broth (HiEncap™ 2X YT Broth)

EC1251CCL

HiEncap™ YT Broth (HiEncap™ 2X YT Broth) is used for the cultivation of recombinant strains of *Escherichia coli*.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	16.000
Yeast extract	10.000
Sodium chloride	5.000
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Each capsule contains 7.75 grams of medium. Suspend 1 capsule in 250 ml (4 capsules in 1000 ml) distilled or purified water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Principle And Interpretation

YT Broth is recommended for use in the cultivation of recombinant strains of *Escherichia coli* (1, 2, 4). It is also used in culture of *E. coli* strains for propagation of M 13 bacteriophages (2,3,4).

These media contain casein enzymic hydrolysate and yeast extract, which supply nitrogenous compounds, vitamin B complex and other essential nutrients and co-factors necessary for the luxuriant growth of recombinant *E. coli* and allows the bacteria to recover from the stress of transformation and grow well. Sodium chloride helps in maintaining isotonic conditions in the medium.

Quality Control

Appearance

Gelatin capsule containing light yellow to beige coloured granular media

Quantity

Each capsule contains 7.75 grams of medium sufficient for 250ml media.

Color and Clarity of Prepared Medium

Light amber coloured clear solution in tubes.

Reaction

Reaction of 3.1% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH

6.80-7.20

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Cultural Response

Organism	Inoculum (CFU)	Growth
Cultural Response		
<i>Escherichia coli</i> ATCC 23724	50-100	good-luxuriant
<i>Escherichia coli</i> ATCC 53868	50-100	good-luxuriant

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Reference

1. Miller H., 1987, Meth. Enzymol; 152, 145.

2. Ausubel F. M., Brent R., Kingston R. E., Moore B. D., Seidman J. G., Smith J. A. and Strohl K., 1994, Current Protocols in Molecular Biology, Vol. I, Current Protocols, New York, N.Y.
3. Davis L. G., Dibner M. D., Battey J. F., 1986, Basic Methods in Molecular Biology, Elsevier, New York, N.Y.
4. Sambrook J., Fritsch E. E. and Maniatis T., 1989, Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.

Revision : 00 / 2014



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.