



Technical Data

Agar Medium C (Sabouraud-Glucose Agar w/ antibiotics)

M1472B

Sabouraud Glucose Agar w/ antibiotics is recommended for selective cultivation of yeasts and moulds in accordance with British Pharmacopoeia 2009.

Composition**

Ingredients	Gms / Litre
Glucose monohydrate	40.000
Peptone (meat and casein)	10.000
Agar	15.000
pH after sterilization (at 25°C)	5.6±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 61.36 grams of dehydrated medium in 995 ml purified/distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. i.e. validated cycle. Aseptically add rehydrated contents of one vial of Tetracycline Selective Supplement (FD196). Mix well before pouring into sterile Petri plates.

Principle And Interpretation

Sabouraud Glucose agar w/ antibiotics is cited as Medium C and recommended for cultivation of yeasts and moulds by British Pharmacopoeia (1). This medium was described originally by Sabouraud (2) for the cultivation of fungi particularly useful for the fungi associated with skin infections. The medium is used with antibiotics such as tetracycline and benzylpenicillin (3) for the isolation of pathogenic fungi from materials containing large numbers of fungi or bacteria.

Peptone (from meat and casein) provides nitrogenous compounds. Glucose monohydrate provides an energy source. Tetracycline and benzyl penicillin inhibits a wide range of Gram-positive and Gram-negative bacteria, which makes the medium selective for fungi (4). The low pH favours fungal growth and inhibits contaminating bacteria from clinical specimens (5). Some pathogenic fungi may produce infective spores, which are easily dispersed in air, so examination should be carried out in safety cabinet.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light amber coloured clear to slightly opalescent gel forms in Petri plates

pH

5.40-5.80

Growth Promotion Test

Growth Promotion was carried out in accordance with the harmonized method of BP, after an incubation at 20-25°C with added Tetracycline Selective Supplement (FD196) for ≤5 days. Recovery rate is considered as 100% for bacteria growth on Soybean Casein Digest Agar and fungus growth on Sabouraud Dextrose Agar

Cultural Response

Organism	Inoculum (CFU)	Growth	Observed Lot value (CFU)	Recovery	Incubation temperature	Incubation period
Cultural Response						

<i>Candida albicans</i> ATCC 10231	50 -100	Luxuriant (white colonies)	25 -100	≥50 %	20 -25 °C	≤5 d
* <i>Aspergillus brasiliensis</i> ATCC 16404	50 -100	luxuriant	25 -100	≥50 %	20 -25 °C	≤5 d
<i>Candida albicans</i> ATCC 2091	50 -100	luxuriant	25 -100	≥50 %	20 -25 °C	≤5 d
<i>Saccharomyces cerevisiae</i> ATCC 9763	50 -100	luxuriant	35 -100	≥50 %	20 -25 °C	≤5 d
<i>Escherichia coli</i> ATCC 25922	≥10 ³	inhibited	0	0 %	20 -25 °C	≤5 d
<i>Escherichia coli</i> ATCC 8739	≥10 ³	inhibited	0	0 %	20 -25 °C	
<i>Escherichia coli</i> NCTC 9002	≥10 ³	inhibited	0	0 %	20 -25 °C	
<i>Trichophyton rubrum</i> ATCC 28191	50-100	good			20 -25 °C	≤5 d
<i>Lactobacillus casei</i> ATCC 334	≥10 ³	inhibited	0	0 %	20 -25 °C	

Key : * - Formerly known as *Aspergillus niger*

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

1. British Pharmacopoeia, 2009, The Stationery Office British Pharmacopoeia
2. Sabouraud K., 1892, Ann. Dermatol. Syphilol, 3:1061.
3. Ajello L., 1957, J. Chron. Dis., 5:545.
4. Lorian (Ed.), 1980, Antibiotics In Laboratory Medicine, Williams and Wilkins, Baltimore.
5. Murray, P. R 2005, In Manual of Clinical Microbiology, 7th ed., ASM, Washington, D.C.

Revision : 1 / 2011



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.