

## UIMC1 Polyclonal Antibody

Catalog number: 13642-1-AP

Size: 33 µg/150 µl


Source: Rabbit

Isotype: IgG


Synonyms:

UIMC1; BRCA1 A complex  
subunit RAP80, RAP80,


Receptor associated protein 80,  
RXRIP110, UIMC1, X2HRIP110




Immunofluorescent analysis of HeLa cells, using UIMC1 antibody 13642-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



mouse colon tissue were subjected to SDS PAGE followed by western blot with 13642-1-AP(UIMC1 antibody) at dilution of 1:300



Immunohistochemistry of paraffin-embedded human testis tissue slide using 13642-1-AP( UIMC1 Antibody) at dilution of 1:50 (under 10x lens)



Immunohistochemistry of

### Applications

Tested applications:

Species specificity:

Calculated UIMC1 MW:

Observed UIMC1 MW:

Positive WB detected in

Positive IP detected in

Positive IHC detected in

Positive IF detected in

Recommended dilution:

ELISA, WB, IF, IHC, IP

Human, Mouse; other species not tested.

719aa,80 kDa

97 kDa,80 kDa,71-75 kDa,61 kDa

Mouse colon tissue, COLO 320 cells, HeLa cells, Raji cells

COLO 320 cells

Human testis tissue

Hela cells, MCF-7 cells

WB: 1:200-1:1000

IP: 1:200-1:2000

IHC: 1:20-1:200

IF: 1:10-1:100

Application key: WB = Western blotting, IHC = Immunohistochemistry, IF = Immunofluorescence, IP = Immunoprecipitation

### Immunogen information

Immunogen:

Ag4582

GenBank accession number:

BC032561

Gene ID (NCBI):

51720

Full name:

Ubiquitin interaction motif containing 1


### Product information

Purification method:

Antigen affinity purification

Storage:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Store at -20°C.



IP Result of anti-UIMC1

paraffin-embedded human testis tissue slide using 13642-1-AP( UIMC1 Antibody) at dilution of 1:50 (under 40x lens)

(IP:13642-1-AP, 4ug;  
Detection:13642-1-AP 1:500)  
with COLO 320 cells lysate  
2400ug.