# X-GAL

Shipping: On Dry/Blue Ice

Catalog number

Batch No.: See vial

BIO-37035: 1 q



Store at -20°C

## Storage and stability:

X-GAL is shipped on dry/blue ice. On arrival store at -20 °C for optimum stability. Repeated freeze/ thaw cycles should be avoided.

#### Expirv:

When stored under the recommended conditions and handled correctly, full activity of the kit is retained until the expiry date on the outer box label.

**Safety precautions:** Harmful if swallowed. Irritating to eyes and skin. Please refer to the material safety data sheet for further information and safety handling practice.



Xn: Harmful

### Quality control specifications:

X-GAL is extensively tested for activity and absence of contamination prior to release.

#### Notes:

Research use only.

# Description

5-bromo-4-chloro-3-indolyl  $\beta$ -D-galactopyranoside (X-GAL) is a chromogenic substrate for  $\beta$ -Galactosidase that forms an intense blue precipitate. It can be used in molecular biology to detect the *gal* gene product, and also in microbiology where it is used to detect microorganisms which have  $\beta$ -Galactosidase activity (usually coliforms). It can be combined with the R-substrates to differentiate between two species of organisms on the same plate. X-GAL is soluble in N, N-dimethylformamide or dimethylsulfoxide (DMSO). Together with appropriate bacterial cloning vectors, host strains, and IPTG, X-GAL provides an easy way to distinguish between positive and negative clones after transformation.

#### **Features**

- Extremely pure, 99.5% by TLC (Thin Layer Chromatography)
- Intense blue precipitate upon hydrolysis

## **Applications**

- Blue/White colony screening to distinguish recombinant (white) and non-recombinant (blue) colonies
- Immunoblotting
- Detect β-galactosidase activity in immunological and histochemical assays
- Microbiology and cell culture media

#### **Protocol**

Prepare a 20 mg/ml stock solution of X-Gal in N,N-dimethylformamide (DMF) or dimethylsulfoxide (DMSO). Sterilization is not required. Store stock solution in glass container protected from light at -20 °C. Solutions may be stored at -20 °C for 6-12 months. If a solution turns pink, it should be discarded. DMF dissolves some plastic materials. Direct addition of DMF containing solution to plastic Petri dishes should be avoided.

## **Product Citations**

- 1. Wilson, A.C., et al. J. Bacteriol. 190(15), 5522-5525(2008).
- 2. Liapis, E., et al. NAR 36(18), 5933-5945 (2008).
- 3. Corbett, D., et al. J. Biol. Chem. 282, 33326-33335 (2007).
- Toledo-Arana, A., et al. J. Bacteriol. 187(15), 5318-5329 (2005).
- 5. Staddon, J.H., et al. Plasmid 56(2), 102-111(2006).
- 6. Maruta, F., et al. Can. Gene Therapy 9(6), 543-552 (2002).
- 7. Marani, M., et al. Mol. Cell. Biol. 22(11), 3577-3589 (2002).

## **Procedures**

Blue/White Colony Screening

- a) X-Gal included in agar: Add 5 ml of X-Gal stock solution and 5 ml of 0.1 M isopropyl-b-D-thiogalactoside (IPTG, BIO-37082) for each 1L of autoclaved media agar (e.g. LB agar) containing appropriate antibiotics just before pouring. The media should be below 55 °C. Plate cells on cooled agar and incubate overnight at 37 °C.
- b) X-Gal applied to top of agar: To a premade LB agar plate (e.g. prepared using LB agar), add 40  $\mu$ l of X-Gal stock solution (at room temperature) and 4  $\mu$ l of a 200 mg/ml solution of IPTG.

Spread solution over the entire surface of the plate. Incubate at 37 °C until the fluid is no longer visible. This may take several hours. Plate cells and incubate overnight at 37 °C.

## **Associated Products**

IPTG Powder	BIO-37036
Quick-Stick Ligase	BIO-27027
SureClean Plus	BIO-37047

Bioline Reagents Ltd UNITED KINGDOM

Bioline USA Inc. USA Bioline GmbH GERMANY Bioline (Aust) Pty. Ltd AUSTRALIA Bioline France FRANCE Meridian Bioscience Asia Pte Ltd SINGAPORE

Tel: +44 (0)20 8830 5300 Fax: +44 (0)20 8452 2822 Tel: +1 508 880 8990 Fax: +1 508 880 8993 Tel: +49 (0)337 168 1229 Fax: +49 (0)3371 68 1244

Tel: +61 (0)2 9209 4180 Fax: +61 (0)2 9209 4763 Tel: +33 (0)1 42 56 04 40 Fax: +33 (0)9 70 06 62 10 Tel: +65 6774 7196 Fax: +65 6774 6441