

Rabbit anti-Reston GP pAb

Catalog #: 0305-001

Lot #: 1306005

Immunogen: Peptide sequence to Reston Ebola virus (RESTV) glycoprotein (GP).

Description: Affinity purified rabbit polyclonal antibody reactive to RESTV GP. The antibody detects recombinant RESTV GP without the transmembrane region (RESTV rGPdTM) in Western blot analysis.

Supplied: 100 μ g of antibody is supplied in PBS at a concentration of 1.92 mg/mL.

Raised in: Rabbits

Purification: Antibody is affinity purified using immobilized immunogen.

Clonality: Polyclonal

Relevance: The antibody can be used for detection of RESTV GP.

Recommended Dilutions:

ELISA: Assay-dependent dilution.

WB: Assay-dependent dilution; internal QC demonstrates detection of recombinant RESTV GPdTM with a use dilution of 50 ng per mL of antibody in diluent using an alkaline phosphatase antibody conjugate and chromogenic substrate for visualization.

Storage: 2-3 weeks +4°C, -20°C long term

Cross Reactivity: No cross reactivity to Sudan ebolavirus (SUDV) GP, Zaire ebolavirus (EBOV) GP, Bundibugyo (BDBV) GP or Marburg (MARV) GP, was observed in Western blot analysis.


Intended for research use only, not for human, therapeutic, or diagnostic applications.

The buyer cannot sell or otherwise transfer this product for Commercial Purposes without written approval of Integrated BioTherapeutics, Inc.

Copyright 2014. Integrated BioTherapeutics, Inc. All rights reserved.

4 Research Court, Suite 300 Rockville, MD 20850 877-411-2041 Services@ibtbioservices.com

Western Blot Data:



50 ng/mL anti-REBOV GP 500 ng/mL anti-REBOV GP

Lane 1: 15 μ L Novex MW markers Lane 2: 15 μ L RESTV rGPdTM supernatant Lane 3: 0.2 μ g EBOV rGPdTM Lane 4: 0.2 μ g SUDV rGPdTM Lane 5: 0.2 μ g MARV rGPdTM Lane 6: 0.2 μ g BDBV rGPdTM Lane 7: 15 μ L Novex MW markers Lane 8: 15 μ L RESTV rGPdTM supernatant Lane 9: 0.2 μ g EBOV rGPdTM Lane 10: 0.2 μ g SUDV rGPdTM Lane 11: 0.2 μ g MARV rGPdTM Lane 12: 0.2 μ g BDBV rGPdTM

Western blots were detected with anti-RESTV GP at 50 ng/mL (left panel) and 500 ng/mL (right panel) and visualized using an anti-rabbit AP conjugate and chromogenic substrate. RESTV GP is visualized as a broad band representing differing glycosylation patterns. Cross reactivity is not detected in the EBOV, SUDV, MARV or BBDV recombinant GP proteins (lanes 3, 4, 5, 6, 9, 10, 11, 12).