



Reinforced Clostridial Broth

M443

Reinforced Clostridial Broth is used for the cultivation and enumeration of *Clostridia* and other anaerobes.

Composition**

| Ingredients | Gms / Litre |
|----------------------------|-------------|
| Casein enzymic hydrolysate | 10.000 |
| Beef extract | 10.000 |
| Yeast extract | 3.000 |
| Dextrose | 5.000 |
| Sodium chloride | 5.000 |
| Starch, soluble | 1.000 |
| L-Cysteine hydrochloride | 0.500 |
| Sodium acetate | 3.000 |
| Agar | 0.500 |
| Final pH (at 25°C) | 6.8±0.2 |

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 38.00 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 10 lbs pressure (115°C) for 15 minutes. Mix well and dispense as desired.

Principle And Interpretation

Reinforced Clostridial Broth is formulated by Hirsch and Grinstead (1). It can be used to initiate growth from small inocula and to obtain the highest viable count of *Clostridia*. Barnes and Ingram used the broth medium for diluting an inoculum of vegetative cells of *Clostridium perfringens* (2). It can be used in studies of spore forming anaerobes, especially *Clostridium butyricum* in cheese, for enumeration of Clostridia in tube dilution counts or for preparation of plates for isolation (3). Other spore forming anaerobes, *Streptococci* and *Lactobacilli* also grow in this media. This is a nonselective enrichment media.

Casein enzymic hydrolysate, yeast extract, beef extract, starch, L-cysteine and sodium acetate provide all the necessary nutrients for the growth of *Clostridia*. Dextrose is a fermentable carbohydrate in the medium while sodium chloride maintains osmotic equilibrium. This media can be made selective by addition of 15-20 mg Polymyxin B per litre of media (1).

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Light yellow coloured clear solution in tubes.

Reaction

Reaction of 3.8% w/v aqueous solution at 25°C . pH : 6.8±0.2

pH

6.60-7.00

Cultural Response

Cultural characteristics observed in an anaerobic atmosphere after an incubation at 35 - 37°C for 24 - 48hours.

Cultural Response

| Organism | Inoculum (CFU) | Growth |
|-------------------------------------------------------------------------|----------------|------------------|
| Cultural Response <i>Clostridium sporogenes</i> ATCC 11437 | 50 - 100 | good - luxuriant |

| | | |
|-----------------------------------------------|---------|------------------|
| <i>Clostridium sporogenes</i> ATCC 19404 | 50 -100 | good - luxuriant |
| <i>Bacteroides vulgatus</i> ATCC 8482 | 50 -100 | good - luxuriant |
| <i>Bacteroides fragilis</i> ATCC 23745 | 50 -100 | good - luxuriant |
| <i>Clostridium perfringenes</i> ATCC 13124 | 50 -100 | good - luxuriant |

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

- 1.Hirsch and Grinsted, 1954, J. Dairy Res., 21:101.
- 2.Barnes and Ingram, 1956, J. Appl. Bact., 19:117.
- 3.Lewis and Angelotti (Eds.), 1964, Examination of Foods for Enteropathogenic and Indicator Bacteria, Dept. of HEW, PHS Publication, 1142, Washington.

Revision : 1 / 2011



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.