



IBT BIOSERVICES

4 Research Court, Suite 300
Rockville, MD 20850
877-411-2041
Services@ibtbioservices.com

Staphylococcus aureus Alpha Hemolysin (Hla)

Catalog #: 1401-002

Lot #: 1601002

Description: *Staphylococcus aureus* alpha hemolysin (Hla), also known as alpha toxin (α -toxin), is expressed in *E. coli* as a recombinant protein without TAG. The theoretical molecular weight of the protein is approximately 33 kDa.

Supplied: 100 μ g is supplied in phosphate buffered saline at a concentration of **0.857 mg/mL**.

Endotoxin: 8.029 Endotoxin units/mg

Storage: -80°C

Relevance: Hla is produced by *S. aureus* and is a major cytotoxic agent. It is expressed as a monomer that binds to the membrane of susceptible cells. Subunits then oligomerize to form heptameric rings with a central pore through which cellular contents leak. The mode of action of *S. aureus* Hla is likely by osmotic lysis.

Recommended Dilutions:


ELISA: Assay-dependent dilution.

WB: Assay-dependent dilution; internal QC demonstrates detection of Hla protein using IBT's mouse anti-Alpha Toxin (6C12) monoclonal antibody (cat# 0210-005) in Western blotting.


Cytotoxicity assay: The protein is active in functional lysis assays with rabbit red blood cells. The protein can be also used in serology assay as coating antigen to detect antibodies against Hla.

For additional *S. aureus* products, please visit:

<http://ibtbioservices.com/index.php/product-and-reagents/staphylococcal-products>



(A) SDS-PAGE of Hla: 1 μ g (lane 1) and 5 μ g (lane 2). (B) Western blot detection of Hla at 5 ng, 10 ng and 50 ng (lanes 1-3), using IBT's mouse anti-Alpha Toxin (6C12) monoclonal antibody at 1.0 μ g/mL and an anti-mouse IgG-HRP conjugate followed by substrate.



Lysis of rabbit red blood cells by *S. aureus* Hla. Red blood cell lysis was determined by absorbance at OD 416 nm after 30 min incubation at 37°C with Hla. Blue circles represents the current lot 1601002 and green circles represents the previous lot 1505002. EC₅₀ was found to be 0.0145 μ g/mL for current lot and 0.0158 μ g/mL for previous lot.

Intended for research use only, not for human, therapeutic, or diagnostic applications.

The buyer cannot sell or otherwise transfer this product for Commercial Purposes without written approval of Integrated BioTherapeutics, Inc.

Copyright 2016. Integrated BioTherapeutics, Inc. All rights reserved.